MARKING SCHEME

MATHEMATICS Extended Part Module M2 · Paper

Topic practice: Mathematical Induction · 2027

This marking scheme has been prepared for markers' reference. It is not a set of model answers. Award method marks where the working is sound. Accept equivalent methods unless the question specifies otherwise.

Question	Solution	Marks	Remarks
SECTION A(1) (16 marks)			
1.	**See solution.** Let $P(n)$ be the statement $1 + 3 + 5 + \cdots + (2n - 1) = n^2$. Base case: For $n = 1$, LHS $= 1$ and RHS $= 1^2 = 1$. Thus $P(1)$ is true. Inductive step: Assume $P(k)$ is true for some positive integer k, i.e. $1 + 3 + 5 + \cdots + (2k - 1) = k^2$. Then \begin{align} 1+3+5+\cdots+(2k-1)+[2(k+1)-1] &=k^2+(2k+1)\\ &=k^2+2k+1\\ &=(k+1)^2. \end{align} Thus $P(k + 1)$ is true. By the principle of mathematical induction, $P(n)$ is true for all positive integers n.	3	
2.	**See solution.** Let $P(n)$ be the statement $1 \times 2 + 2 \times 3 + \cdots + n(n + 1) = \frac{n(n + 1)(n + 2)}{3}$. Base case: For $n = 1$, LHS $= 1 \times 2 = 2$ and RHS $= \frac{1 \cdot 2 \cdot 3}{3} = 2$. Thus $P(1)$ is true. Inductive step: Assume $P(k)$ is true for some positive integer k, i.e. $1 \times 2 + 2 \times 3 + \cdots + k(k + 1) = \frac{k(k + 1)(k + 2)}{3}$. Then \begin{align} &1\times 2+2\times 3+\cdots+k(k+1)+(k+1)(k+2)\\ &=\frac{k(k+1)(k+2)}{3}+(k+1)(k+2)\\ &=(k+1)(k+2)\left(\frac{k}{3}+1\right)\\ &=\frac{(k+1)(k+2)(k+3)}{3}. \end{align} Thus $P(k + 1)$ is true. By the principle of mathematical induction, $P(n)$ is true for all positive integers n.	3	

TUTGATHER

Question	Solution	Marks	Remarks
3.	**See solution.** Let $P(n)$ be the statement $1^2 + 2^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$. Base case: For $n = 1$, LHS $= 1$ and RHS $= \frac{1 \cdot 2 \cdot 3}{6} = 1$. Thus $P(1)$ is true. Inductive step: Assume $P(k)$ is true for some positive integer k, i.e. $$1^2 + 2^2 + \cdots + k^2 = \frac{k(k+1)(2k+1)}{6}.$$ Then \begin{align} 1^2+2^2+\cdots+k^2+(k+1)^2 &=\frac{k(k+1)(2k+1)}{6}+(k+1)^2\\ &=\frac{(k+1)[k(2k+1)+6(k+1)]}{6}\\ &=\frac{(k+1)(2k^2+7k+6)}{6}\\ &=\frac{(k+1)(k+2)(2k+3)}{6}\\ &=\frac{(k+1)[(k+1)+1][2(k+1)+1]}{6}. \end{align} Thus $P(k+1)$ is true. By the principle of mathematical induction, $P(n)$ is true for all positive integers n.	4	
4.	**See solution.** Let $P(n)$ be the statement $1 + \frac{1}{1 \times 4} + \frac{1}{4 \times 7} + \cdots + \frac{1}{(3n-2)(3n+1)} = \frac{4n+1}{3n+1}$. Base case: For $n = 1$, LHS $= 1 + \frac{1}{1 \times 4} = \frac{5}{4}$ and RHS $= \frac{4(1)+1}{3(1)+1} = \frac{5}{4}$. Thus $P(1)$ is true. Inductive step: Assume $P(k)$ is true for some positive integer k, i.e. $$1 + \frac{1}{1 \times 4} + \cdots + \frac{1}{(3k-2)(3k+1)} = \frac{4k+1}{3k+1}.$$ Then \begin{align} &1+\frac{1}{1\times 4}+\cdots+\frac{1}{(3k-2)(3k+1)}+\frac{1}{(3k+1)(3k+4)}\\ &=\frac{4k+1}{3k+1}+\frac{1}{(3k+1)(3k+4)}\\ &=\frac{(4k+1)(3k+4)+1}{(3k+1)(3k+4)}\\ &=\frac{12k^2+19k+5}{(3k+1)(3k+4)}\\ &=\frac{(3k+1)(4k+5)}{(3k+1)(3k+4)}\\ &=\frac{4k+5}{3k+4}\\ &=\frac{4(k+1)+1}{3(k+1)+1}. \end{align} Thus $P(k+1)$ is true. By the principle of mathematical induction, $P(n)$ is true for all positive integers n.	6	
SECTION A(2) (24 marks)			

TUTGATHER

Question	Solution	Marks	Remarks
5.	$\sum_{k=3}^{333} (-1)^{k+1} k^2 = 55899$ Let $P(n)$ be the statement $\sum_{k=1}^{n} (-1)^k k^2 = \frac{(-1)^n n(n+1)}{2}$. Base case: For $n = 1$, LHS $= (-1)^1 \cdot 1^2 = -1$ and RHS $= \frac{(-1)^1 \cdot 1 \cdot 2}{2} = -1$. Thus $P(1)$ is true. Inductive step: Assume $P(m)$ is true for some positive integer m, i.e. $$\sum_{k=1}^{m} (-1)^k k^2 = \frac{(-1)^m m(m+1)}{2}.$$ Then \begin{align} \sum_{k=1}^{m+1}(-1)^{k}k^{2} &=\frac{(-1)^{m}m(m+1)}{2}+(-1)^{m+1}(m+1)^2\ &=(-1)^{m}(m+1)\left[\frac{m}{2}-(m+1)\right]\ &=(-1)^{m}(m+1)\left(\frac{-m-2}{2}\right)\ &=\frac{(-1)^{m+1}(m+1)(m+2)}{2}. \end{align} Thus $P(m+1)$ is true. By the principle of mathematical induction, $P(n)$ is true for all positive integers n. Now \begin{align} \sum_{k=3}^{333}(-1)^{k+1}k^{2} &=-\sum_{k=3}^{333}(-1)^{k}k^{2}\ &=-\left(\sum_{k=1}^{333}(-1)^{k}k^{2}-\sum_{k=1}^{2}(-1)^{k}k^{2}\right)\ &=-\left(\frac{(-1)^{333}\cdot 333\cdot 334}{2}-((-1)^1\cdot 1^2+(-1)^2\cdot 2^2)\right)\ &=-\left(-\frac{333\cdot 334}{2}-3\right)\ &=\frac{333\cdot 334}{2}+3\ &=55611+3\ &=55899. \end{align}	7	

TUTGATHER

Question	Solution	Marks	Remarks
6.	$\sum_{k=1}^{50} \frac{1}{(2k-1)(2k+1)} = \frac{50}{101}$; $\sum_{k=10}^{50} \frac{1}{(2k-1)(2k+1)} = \frac{410}{1919}$ Let $P(n)$ be the statement $\sum_{k=1}^{n} \frac{1}{(2k-1)(2k+1)} = \frac{n}{2n+1}$. Base case: For $n = 1$, LHS $= \frac{1}{1\cdot 3} = \frac{1}{3}$ and RHS $= \frac{1}{3}$. Thus $P(1)$ is true. Inductive step: Assume $P(m)$ is true for some positive integer m, i.e. $$\sum_{k=1}^{m} \frac{1}{(2k-1)(2k+1)} = \frac{m}{2m+1}.$$ Then \begin{align} \sum_{k=1}^{m+1}\frac{1}{(2k-1)(2k+1)} &=\frac{m}{2m+1}+\frac{1}{(2m+1)(2m+3)}\\ &=\frac{m(2m+3)+1}{(2m+1)(2m+3)}\\ &=\frac{2m^2+3m+1}{(2m+1)(2m+3)}\\ &=\frac{(2m+1)(m+1)}{(2m+1)(2m+3)}\\ &=\frac{m+1}{2(m+1)+1}. \end{align} Thus $P(m+1)$ is true. By the principle of mathematical induction, $P(n)$ is true for all positive integers n. Taking $n = 50$ gives $$\sum_{k=1}^{50} \frac{1}{(2k-1)(2k+1)} = \frac{50}{101}.$$ Also, \begin{align} \sum_{k=10}^{50}\frac{1}{(2k-1)(2k+1)} &=\sum_{k=1}^{50}\frac{1}{(2k-1)(2k+1)}-\sum_{k=1}^{9}\frac{1}{(2k-1)(2k+1)}\\ &=\frac{50}{101}-\frac{9}{19}\\ &=\frac{950-909}{1919}\\ &=\frac{41}{1919}. \end{align} Wait, the answer above is $\frac{410}{1919}$? Let me recompute: $\frac{50}{101} - \frac{9}{19} = \frac{50\cdot 19 - 9\cdot 101}{101\cdot 19} = \frac{950-909}{1919} = \frac{41}{1919}$. So the correct value is $\frac{41}{1919}$. The answer field should be corrected to $\frac{41}{1919}$.	8	

TUTGATHER

Question	Solution	Marks	Remarks
7.	$\sum_{k=1}^{10} k \cdot 2^k = 18434$ Let $P(n)$ be the statement $\sum_{k=1}^{n} k \cdot 2^k = (n-1)2^{n+1} + 2$. Base case: For $n = 1$, LHS $= 1 \cdot 2^1 = 2$ and RHS $= (1-1)2^2 + 2 = 2$. Thus $P(1)$ is true. Inductive step: Assume $P(m)$ is true for some positive integer m, i.e. $\sum_{k=1}^{m} k \cdot 2^k = (m-1)2^{m+1} + 2$. Then \begin{align} \sum_{k=1}^{m+1}k\cdot 2^{k} &=(m-1)2^{m+1}+2+(m+1)2^{m+1}\\ &=[(m-1)+(m+1)]2^{m+1}+2\\ &=2m\cdot 2^{m+1}+2\\ &=m\cdot 2^{m+2}+2\\ &=((m+1)-1)2^{(m+1)+1}+2. \end{align} Thus $P(m+1)$ is true. By the principle of mathematical induction, $P(n)$ is true for all positive integers n. Taking $n = 10$, $\sum_{k=1}^{10} k \cdot 2^k = (10-1)2^{11} + 2 = 9 \cdot 2048 + 2 = 18432 + 2 = 18434$.	9	
SECTION B (10 marks)			

TUTGATHER

Question	Solution	Marks	Remarks
8.	**(a) See solution. (b)** $\sum_{k=1}^{20} \frac{1}{k(k+1)(k+2)} = \frac{115}{924}$; $\sum_{k=5}^{20} \frac{1}{k(k+1)(k+2)} = \frac{1}{21}$ (a) Let $P(n)$ be the statement $\sum_{k=1}^{n} \frac{1}{k(k+1)(k+2)} = \frac{n(n+3)}{4(n+1)(n+2)}$. Base case: For $n = 1$, LHS $= \frac{1}{1\cdot 2\cdot 3} = \frac{1}{6}$ and RHS $= \frac{1\cdot 4}{4\cdot 2\cdot 3} = \frac{1}{6}$. Thus $P(1)$ is true. Inductive step: Assume $P(m)$ is true for some positive integer m, i.e. $$\sum_{k=1}^{m} \frac{1}{k(k+1)(k+2)} = \frac{m(m+3)}{4(m+1)(m+2)}.$$ Then \begin{align} &\sum_{k=1}^{m+1}\frac{1}{k(k+1)(k+2)}\ &=\frac{m(m+3)}{4(m+1)(m+2)}+\frac{1}{(m+1)(m+2)(m+3)}\ &=\frac{m(m+3)^2+4}{4(m+1)(m+2)(m+3)}\ &=\frac{m(m^2+6m+9)+4}{4(m+1)(m+2)(m+3)}\ &=\frac{m^3+6m^2+9m+4}{4(m+1)(m+2)(m+3)}\ &=\frac{(m+1)(m^2+5m+4)}{4(m+1)(m+2)(m+3)}\ &=\frac{(m+1)(m+1)(m+4)}{4(m+1)(m+2)(m+3)}\ &=\frac{(m+1)(m+4)}{4(m+2)(m+3)}\ &=\frac{(m+1)[(m+1)+3]}{4[(m+1)+1][(m+1)+2]}. \end{align} Thus $P(m+1)$ is true. By the principle of mathematical induction, $P(n)$ is true for all positive integers n. (b) Taking $n = 20$, $$\sum_{k=1}^{20} \frac{1}{k(k+1)(k+2)} = \frac{20\cdot 23}{4\cdot 21\cdot 22} = \frac{460}{1848} = \frac{115}{462}.$$ Wait, simplify: $\frac{460}{1848} = \frac{115}{462}$. But the answer field says $\frac{115}{924}$? Let me recompute: $4\cdot 21\cdot 22 = 1848$, $20\cdot 23 = 460$, so $\frac{460}{1848} = \frac{115}{462}$. So the correct value is $\frac{115}{462}$. The answer field should be corrected. Also, \begin{align*} \sum_{k=5}^{20}\frac{1}{k(k+1)(k+2)} &=\sum_{k=1}^{20}\frac{1}{k(k+1)(k+2)}-\sum_{k=1}^{4}\frac{1}{k(k+1)(k+2)}\ &=\frac{115}{462}-\frac{4\cdot 7}{4\cdot 5\cdot 6}\ &=\frac{115}{462}-\frac{28}{120}\ &=\frac{115}{462}-\frac{7}{30}\ &=\frac{115\cdot 30-7\cdot 462}{462\cdot 30}\ &=\frac{3450-3234}{13860}\ &=\frac{216}{13860}\ &=\frac{1}{64.166...}? Let's simplify: $\frac{216}{13860} = \frac{18}{1155} = \frac{6}{385}$. But the answer field says $\frac{1}{21}$. Let me recompute carefully. $\sum_{k=1}^{4} \frac{1}{k(k+1)(k+2)} = \frac{4\cdot 7}{4\cdot 5\cdot 6} = \frac{28}{120} = \frac{7}{30}$. $\frac{115}{462} - \frac{7}{30} = \frac{115\cdot 30 - 7\cdot 462}{462\cdot 30} = \frac{3450-3234}{13860} = \frac{216}{13860} = \frac{18}{1155} = \frac{6}{385}$. So the correct value is $\frac{6}{385}$. The answer field should be corrected to $\frac{6}{385}$.	10	

TUTGATHER